



Cytotoxic Effects of Hydroxy Coumarin Derivatives on Mouse Neuroblastoma N2a Cell Line

Javad Sargolzaei^{a,c}, Hamid Sadeghian^b, Sanaz Golahmadi^c, Mohammad Soukhtanloo^{d,e*}

^aDepartment of Biology, Faculty of Science, Arak University, Arak, Iran, ^bDepartment of laboratory Sciences, School of Paramedical Sciences, Mashhad University of Medical Sciences, Mashhad, Iran, ^cDepartment of Biochemistry, Payame Noor University, Mashhad, Iran, ^dDepartment of Clinical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran, ^ePharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad 917794-8564, Iran.

Abstract

Neuroblastoma is one of the nervous system cancers, which approximately consists of 9% of childhood cancers. In this study, we evaluated the toxic effects of prenyl hydroxy coumarin derivatives on apoptosis of the neuroblastoma cell line N2A. N2A cells were cultured in DMEM medium, then the effects of different concentrations (0.75–200 µg/mL) of prenyl hydroxy coumarin derivatives during 24, 48, and 72 h were studied. Cell viability was quantified by MTT assay; apoptotic cells were determined using PI staining of DNA fragmentation through flow cytometry (sub-G1 peak). The toxic effect of 3- farnesyl oxi coumarin in the N2A cell starts at 6.25 µg/ml and increases relatively depending on rising in concentration and over time. The toxicity and apoptosis of 3- farnesyl and 6- farnesyl oxi coumarin is more than 3- Geranyl and 6-Geranyl oxi coumarin. Prenyl hydroxy coumarin induces peak sub-G1 in flow cytometry compared to the control group, indicating that induced toxicity, which is involved in apoptotic cell death. Different concentrations of derivatives (0.75-200 µg/mL) in lymphocytes did not induce any anti-proliferative effect in 24 h. In conclusion, prenyl hydroxy coumarin derivatives induce apoptotic effects in the N2A cell line. Then prenyl hydroxy coumarin derivatives sound to be chemotherapeutic agents for the neuroblastoma cancer cells.

Keywords: Apoptosis, N2A cell line, Prenyl hydroxy coumarin derivatives.

1. Introduction

Cancer is currently one of the leading causes of death in the world, with around 14 million new cases in 2012 and approximately 9.6 million deaths worldwide [1]. There will

be 21.7 million new cancer cases and about 13.1 million cancer deaths in 2030 [2]. Anticancer drugs prevent the uncontrolled division of cancer cells, through DNA intercalating, cross-linking of DNA cancer

Corresponding Authors: Mohammad Soukhtanloo, Department of Clinical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran; Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad 917794-8564, Iran

Tel: (+98)5138002363

Email: soukhtanloom@mums.ac.ir

Cite this article as: Sargolzaei J, Sadeghian H, Golahmadi S, Soukhtanloo M, Cytotoxic Effects of Hydroxy Coumarin Derivatives on Mouse Neuroblastoma N2a Cell Line, 2020, 16 (3): 95-106.

cells, inhibition of DNA topoisomerase, and disruptors in the formation of cytoskeleton cancer cells [3]. Breast cancer is one of the major and deadly cancers, which is the second most common cancer in women and also the fourth most common cause of mortality and morbidity [4]. NF- κ B is a transcription factor that inhibits the activation of apoptosis, which is inhibited by (S)-ginsenoside Rg3 (SRg3) in the MDA-MB-231 cell line [5]. Another study on the combination of Bacopaside (bac) I and II on the MDA-MB-231 class induced apoptosis and inhibited proliferation [6].

Neuroblastoma is one of the most common cancers in children. It is a neuroendocrine tumor, often originating from the sympathetic nervous system (SNS) [7]. About 50% of these tumors appear in the adrenal medulla, and the remaining 50% are in the paraspinal sympathetic ganglia region of the neck, chest, abdomen, and pelvis [8]. Neuroblastoma is one of the most common types of cancers in childhood, accounting for about 7-10 percent of all childhood cancers, and averaging about one per 7,000 live births each year, so about 700 new cases are reported in the United States each year [8-10].

The prevalence of neuroblastoma is uniform throughout the world (at least for western societies); the median age of diagnosis is 18 months [8, 11]. The etiology of this disease is still unknown, but about 40% at 12 months old and about 75% and 98% are diagnosed at 4 and 10 years, respectively, which are probably not affected by the environment [11].

Apoptosis is a physiological cell death that is essential for the evolution and homeostasis of the tissues [12]. Therefore, apoptotic pathways can be therapeutically exploited for cancer treatment and regulated by the balance between pro- and anti-apoptotic ones [13]. Some anticancer drugs induce apoptotic pathways in cancer cell [14]. Anticancer drugs have many characteristics that include: first, having high efficacy at usage in targeted therapies (drugs that specifically inhibit the growth of cancer cells); second, they have low toxicity for body organs [15]. Antioxidants for a long time have been effective agents in the prevention and medication of many diseases, including cancer. Therefore, many fruits and vegetables are sources of abundant antioxidants [16, 17]. Coumarin is a phenolic compound, which has a double-ring that consists of benzene and α -pyrones [18]. It naturally contains the extract of cassia leaf oil, lavender oil, Rutaceae, and Umbelliferae [18, 19]. Interestingly, coumarin also exists in microorganisms, such as *Streptomyces* and *Aspergillus* species [20, 21].

Coumarin has extensive effects on the body tissues, including different biological effects: anti-inflammatory, antimicrobial,

antioxidant, anti-tumor, anti-influenza, and antihyperlipidemic activity [22-27].

In this study, we evaluate the anti-proliferative effects of complete series of prenyl hydroxy coumarin derivatives such as 3-farnesyl oxi coumarin, 3-geranyl oxi coumarin, 6-farnesyl oxi coumarin, and 6-geranyl oxi coumarin on the mouse neuroblastoma N2A cell line and human lymphocytes [28].

2. Materials and Methods

2.1. Chemicals

Sodium citrate, 3-(4,5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium (MTT), Trypan blue, Triton X-100, and the fluorescent probe propidium iodide (PI) were purchased from Sigma-Aldrich (St. Louis, MO, USA). DMEM and FCS were purchased from Gibco (Grand Island, NY, USA).

2.2. Cell culture and Experiments

All experiments in this study were performed on mice neuroblastoma N2A cell lines (tumor-like neuroblasts). This cell lines were provided by the National Cell Bank of Iran (NCBI, Pasteur Institute of Iran, Tehran) and cultured in Dulbecco Modified Eagle's Medium (DMEM). The N2A cells were cultured in DMEM media with FBS 10% and Penicillin-Streptomycin solution (100 unit/ml). The cells were maintained to 95% air and 5% CO₂ at 37°C in a humidified incubator. The cells were passaged every 3-4 days with trypsin EDTA in a flask.

Human lymphocytes were isolated from heparinized blood of healthy volunteers by

Ficoll-Hypaque density gradient centrifugation. They were suspended at a density of 1.0×10^6 cells/ml in a complete DMEM medium and cultured, as described previously[29].

2.3. Cell Viability

The MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) assay is used to determine cell viability[30]. In brief, 10000 cell/well were seeded into flat-bottom 96-well cell culture plates and were incubated overnight into 95% air and 5%CO₂ at 37°C in a humidified incubator. Then cells were treated with different concentrations (0.75–200 µg/mL) of 3-Farnesyl oxi coumarin, 3-Geranyl oxi coumarin, 6-Farnesyl oxi coumarin and 6-Geranyl oxi coumarin for 24, 48 and 72 hours. After incubation, the supernatant was removed and, MTT (5 mg/ml in PBS) was added (the plate was wrapped with aluminum foil to prevent light exposure) then it was kept in the incubator for 4 hours. In the final step, 100 µl of DMSO was added and incubated for 3-5 minutes with gentle shaking (we made sure the crystals entirely dissolved). The absorbance was measured at 570 nm by ELISA reader (Stat fox).

2.4. Staining DNA Using Propidium Iodide

Apoptotic cells were detected using PI staining of cells by flow cytometry to detect the so-called sub-G1 peak[31]. Briefly, N2A cells were seeded 10^5 cell/well into a 24-well plate and allowed them to grow overnight. They were treated with different concentrations of (0.75–200 µg/mL) 3-farnesyl

oxi coumarin, 3-geranyl oxi coumarin, 6-farnesyl oxi coumarin and 6-geranyl oxi coumarin for 48 hours. The floating and adherent cells were harvested and incubated at 4°C overnight in the dark place using 750 µL of a hypotonic buffer containing (50 µg/mL PI in 0.1% sodium citrate plus 0.1% Triton X-100). The number of cells in the staining solution should be at a constant ratio of 1×10^6 cells per ml of hypotonic PI staining solution. Then they should be analyzed by a BD FACSCalibur™ flow cytometer (Becton Dickinson, Mountain View, CA, USA). About 10,000 events were acquired with FACS [32].

2.5. Statistical Analysis

All data are presented as Mean \pm SEM from 3 experiments. One-way analysis of variance (ANOVA), followed by the Bonferroni post-test for multiple-group comparisons were used for data analysis. A p-value of less than 0.05 was considered statistically significant.

3. Results and Discussion

3.1. Effects of Hydroxy Coumarin Derivatives on Cell Proliferation and Viability of Tumor Cells

Prenyl hydroxy coumarin derivatives including 3-Farnesyl oxi coumarin, 3-Geranyl




Figure 1. Cell survival rate of N2A cells evaluated by MTT. The cells at a density of 10^6 cells/mL were cultured for 24 h, 48h and 72h in the absence and presence of different concentrations of Prenyl hydroxy coumarin derivatives (3-Farnesyl oxi coumarin, 3-Geranyl oxi coumarin, 6-Farnesyl oxi coumarin and 6-Geranyl oxi coumarin) (0–200 µg/mL). Data are presented as means \pm SD of three independent experiments. Profiles A, B and C are 24h, 48 and 72h, respectively. *p < 0.05, **p < 0.01, and ***p < 0.001 compared to control.

oxi coumarin, 6-Farnesyl oxi coumarin, and 6-Geranyl oxi coumarin (0.75–200 µg/mL) have dose-dependent cytotoxic effects on neuroblastoma cells (Figure 1). Therefore, the coumarin derivatives have enhanced concentration-dependent cytotoxic effects that the farnesyl hydroxy coumarin derivatives (3-Farnesyl oxi coumarin and 6-Farnesyl oxi coumarin) and the geranyl hydroxy coumarin derivatives (3-Geranyl oxi coumarin and 6-Geranyl oxi coumarin) exhibit the most anti-proliferative properties in 24 hours and 72 hours, respectively (Figure 1: A and C). Also, there is a significant difference between the neuroblastoma cells treated with Prenyl hydroxy coumarin derivatives (The IC₅₀ values of these derivatives against N2A cell lines after 24 h, 48h and 72h were determined (shown in Table 1, after 24, 48 and 72 hours (P <0.001)).

cells when compared to the control, thus suggesting that apoptotic cell death is involved in its toxicity. IC₅₀ values which were obtained from flow cytometry histogram of 6-Farnesyl oxi coumarin, 6- Geranyl oxi coumarin, 3-Farnesyl oxi coumarin and 3-Geranyl oxi coumarin in 48h were 35.8 µg/ml, 47.5 µg/ml, 98.4 µg/ml, and 197.2 µg/ml, respectively. Thus toxicity and apoptosis in compounds such as 6- Farnesyl oxi coumarin and 6-Geranyl oxi coumarin are more than 3-Farnesyl oxi coumarin and 3-Geranyl oxi coumarin. Results indicate a prenyl hydroxy coumarin derivative induced apoptosis in a concentration-dependent manner (Figure 2). As shown in Fig. 2, the Prenyl hydroxy coumarin derivative induces and also loses the membrane integrity of the cell at various concentrations of these Prenyl hydroxy coumarin derivatives in 48 hours.

Table 1. Doses inducing IC₅₀ of Prenyl hydroxy coumarin derivatives against N2A cell lines.

Cell line	Compound/drugs	24h	48h	72h
N2A	3-Farnesyl oxi coumarin (µg/ml)	12.6±2.1	82.8±1.9	188.4±2.6
	6-Farnesyl oxi coumarin (µg/ml)	9.25±1.8	37.5±1.2	83.6±1.9
	3-Geranyl oxi coumarin (µg/ml)	83.2±3.2	132.5±1.6	128.2±2.2
	6-Geranyl oxi coumarin (µg/ml)	68.9±1.8	25±1.1	95.6±1.8

The cells incubated with different concentration of Prenyl hydroxy coumarin derivatives for 24 h, 48h and 72h. IC₅₀ values were expressed as the mean ± SD (n = 3)

3.2. Effects of Prenyl Hydroxy Coumarin Derivatives on Apoptosis in Neuroblastoma Cells

Prenyl hydroxy coumarin derivatives (IC₅₀, low, and more concentration) induce a sub-G1 peak in the flowcytometry histogram of treated

3.3. Effects of Hydroxy Coumarin Derivatives in Normal Cells

Natural coumarin and coumarin derivatives as anticancer agents have been investigated by many researchers [33-37]. Coumarins have excellent properties, such as the ease of




Figure 2. Flow cytometry profiles of the cells 48 h after Prenyl hydroxy coumarin derivatives exposure, stained with PI. Data are means \pm SD of three experiments. Cells were treated with IC_{50} , low and more concentration of Prenyl hydroxy coumarin derivatives for 48 h. Flow cytometry histograms analysis of the cells stained with PI and analyzed by flow cytometry assay. The percent of the cells in Sub-G1 peak were analyzed by *FlowJo* software. Profiles (A–D) are 6-Farnesyl oxi coumarin, 6-Geranyl oxi coumarin, 3-Farnesyl oxi coumarin, and 3-Geranyl oxi coumarin, respectively. *** $P < 0.001$ versus control cells (0 $\mu\text{g/mL}$) regarding Prenyl hydroxy coumarin derivatives; ** $P < 0.01$ versus control cells (0 $\mu\text{g/mL}$) regarding Prenyl hydroxy coumarin derivatives.

chemical modification on them, as compared to the rest of the compounds, the richness of many plants of coumarin, and low toxicity[38]. Hydroxy coumarin derivatives participate in the proliferation process of human lymphocytes. To determine whether the induction of Hydroxy coumarin derivatives was relevant to the potentiation effect, experiments were performed in human lymphocytes. In lymphocytes, exposure to 0.75-200 $\mu\text{g/mL}$ Hydroxy coumarin derivatives (200 $\mu\text{g/mL}$ maximum concentration achievable) did not induce any antiproliferative effect (survival fraction $> 98\%$) in 24 hours. There was no effect on lymphocytes, which were incubated for 24 hours with Hydroxy coumarin derivatives. Then lymphocytes were incubated for 48 h and 72 h with 0.75-200 $\mu\text{g/mL}$ Hydroxy coumarin

derivatives. In the maximum concentration, a small potentiation effect was seen (survival fraction $> 98\%$ for all fractions, except for 6-Farnesyl oxi coumarin). The survival fraction for 6-Farnesyl oxi coumarin in 48h and 72h with 200 $\mu\text{g/mL}$ was $\sim 75.5\%$, and 72.2% , respectively, and all concentration, failed to induce cytotoxicity (survival fraction $\sim 97\%$). The effect of Hydroxy coumarin derivatives on lymphocytes as normal cells is presented in Fig. 3.

Coumarins, depending on the pattern of substitution and replacement of functional groups on the coumarin ring, act differently on different cancer cells[39]. Isoprenylated coumarin derivatives with systematic variations in the tail length (5, 10, or 15 carbons) and substitution position on the coumarin scaffold (at 3-, 6- or 7-position),




Figure 3. Cell survival rate of human lymphocyte evaluated by MTT. The cells at a density of 10^6 cells/mL were cultured for 24 h, 48h and 72h in presence of different concentrations of Prenyl hydroxy coumarin derivatives (3-Farnesyl oxi coumarin , 3-Geranyl oxi coumarin, 6-Farnesyl oxi coumarin and 6-Geranyl oxi coumarin) (0–200 µg/mL). Data are presented as Mean \pm SD of three independent experiments. After 48, 72 hour in 200 µg/mL concentration viability decreased. Profiles A, B and C are 24h, 48 and 72h respectively. ** $P < 0.01$ versus control cells (0 µg/mL) regarding Prenyl hydroxy coumarin derivatives.

causing synthesized coumarin derivatives of the coumarin [40]. According to this study, 6-substituted farnesylated ether coumarin derivative has the highest cytotoxic activity in the 24-hour incubation among the isoprenylated derivatives of coumarin[41].

In comparing all of them, Farnesyl oxi coumarin is stronger than Geranyl oxi coumarin and has significant biological activity. This study investigates the cytotoxic and proapoptotic effects of Prenyl hydroxy

coumarin derivatives on neuroblastoma cells (N2A). In the author's opinion, this is the first report of the toxicity of Prenyl hydroxy coumarin derivatives (3-Farnesyl oxi coumarin, 3-Geranyl oxi coumarin, 6-Farnesyl oxi coumarin, and 6-Geranyl oxi coumarin) in this cancer cell line. Our data confirm that the Prenyl hydroxy coumarin derivative has cytotoxic effects against neuroblastoma cells, which is consistent with previous studies of other derivatives of coumarin, such as Furano

coumarins[41, 42], Dihydrofurano coumarins [43], Phenyl coumarins [44], and Bicomarins [45]. Also, our data show that Prenyl hydroxy coumarin derivatives constitute a significant inhibitor of the proliferation of neuroblastoma cells (N2A).

It was found that among the Prenyl hydroxy coumarin derivatives, 3-Farnesyl oxi coumarin, and 6-Farnesyl oxi coumarin, have the most anti-proliferative activity after 24 h and 48 h, respectively. Previous results of other coumarin derivatives indicate that apoptosis is involved in the death of induced cancer cells [46]. Also, Hydroxy coumarin derivations treatment did not affect the proliferation of normal lymphocytic cells, indicating a degree of specificity for N2A cells.

Coumarin and its derivatives show cytotoxic effects in dose-dependent behaviors [48]. Cancer cells treated by coumarin and its derivatives show apoptotic properties such as nucleoplasm and cytoplasmic condensation, DNA fragmentation, formation membrane-bound apoptotic bodies, and loss of microvilli[48, 49].

The neuroblastoma cells were treated with different concentrations of Prenyl hydroxy coumarin derivatives (IC_{50} , low and more concentration) with a hypotonic phosphate–citrate buffer. The reported results indicate that after incubation of the neuroblastoma cells, DNA fragmentation, and small apoptotic components were observed. At present, the rate of apoptosis and DNA fragmentation is determined using PI staining of DNA fragmentation with flow cytometry (sub-G1

peak). In this staining with PI, the cells that have lost their DNA are less colored and appear to the left of the peak of the G1 peak[47-50]. This study shows that the Prenyl hydroxy coumarin derivatives prevent the proliferation of N2A tumor cells by interfering with apoptosis. These derivatives can be considered as promising chemotherapy agents for cancer treatment. At the end of 24h, 48h, and 72h incubation, Hydroxy coumarin derivatives did not show any cytotoxic effect on lymphocytes (Figure 3: A , B and C).

4. Conclusion

Neuroblastomas are cancers that start in early nerve cells (called neuroblasts) of the sympathetic nervous system, and they consist of 9% of childhood cancers .Apoptosis which is happening during development and aging is a homeostatic mechanism to preserve cell populations in tissues. Some anticancer drugs induce apoptotic pathways in cancer cell [14, 15]. Hydroxy coumarin derivatives have eligible significant anti-proliferative effects on neuroblastoma, renal, lung, and breast cancers, also leukemia[39, 51]. Notably, coumarin and its derivatives inhibit the growth of cancer cells, yet they do not affect normal cells [39, 48]. Therefore, these derivatives are more effective in cells whose growth or signaling pathways have changed, rather than normal cells. In the present study, five derivatives of Prenyl hydroxy coumarin were used in different concentration on N2a cell line. The results confirmed that 3-Farnesyl oxi coumarin and

6-Farnesyl oxo coumarin have more anti-proliferative and anti-apoptotic effects than 3-Geranyl oxo coumarin and 6-Geranyl oxo coumarin after 24 h and 48 h incubation with N2a cell line. These data were validated in high and low concentrations of IC50 through flow cytometry. These derivatives had no anti-proliferative and anti-apoptotic activities on normal lymphocytic cells. Then, findings indicated that these derivatives are important as anti-cancer agents. However, the precise molecular mechanisms of these Prenyl hydroxy coumarin derivatives are still unclear, but there is an ongoing research, that will soon provide a systematic mechanism of these derivatives.

References

- [1] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA. Cancer J. Clin.* (2018) 68 (6): 394-424.
- [2] Fitzmaurice C, Allen C, Barber RM, Barregard L, Bhutta ZA, Brenner H, Dicker DJ, Chimed-Orchir O, Dandona R, Dandona L and Fleming T. Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 32 cancer groups, 1990 to 2015: a systematic analysis for the global burden of disease study. *J.A.M.A. Oncol.* (2015) 3(4): 524-548.
- [3] Kumar N, Bhatnagar A and Dudhe R. Synthesis of 3-(4, 5-dihydro-1-phenyl-5-substituted phenyl-1H-pyrazol-3-yl)-2H-chromen-2-one derivatives and evaluation of their anticancer activity. *Arab. J. Chem.* (2017) S2443-S2452.
- [4] Palethorpe HM, Smith E, Tomita Y, Nakhjavani M, Yool AJ, Price TJ, Young JP, Townsend AR and Hardingham JE. Bacopasides I and II Act in Synergy to Inhibit the Growth, Migration and Invasion of Breast Cancer Cell Lines. *Molecules* (2019) 24 (19): 3539.
- [5] Nakhjavani M, Hardingham JE, Palethorpe HM, Tomita Y, Smith E, Price TJ and Townsend AR. Ginsenoside Rg3: Potential molecular targets and therapeutic indication in metastatic breast cancer. *Medicines* (2019) 6 (1): 17.
- [6] Nakhjavani M, Palethorpe HM, Tomita Y, Smith E, Price TJ, Yool AJ, Pei JV, Townsend AR and Hardingham JE. Stereoselective Anti-Cancer Activities of Ginsenoside Rg3 on Triple Negative Breast Cancer Cell Models. *Pharmaceuticals* (2019) 12 (3): 117.
- [7] Louis CU and Shohet JM. Neuroblastoma: molecular pathogenesis and therapy. *Annu. Rev. Med.* (2015) 66: 49-63.
- [8] Gurney JG, Ross JA, Wall DA and Bleyer WA, Severson RK and Robison LL. Infant cancer in the US: histology-specific incidence and trends, 1973 to 1992. *J. Pediatr. Hematol. Oncol.* (1997) 19 (5): 428-432.
- [9] Pizzo PA and Poplack DG. *Principles and practice of pediatric oncology: Lippincott Williams & Wilkins: USA* (2015).
- [10] Tolbert VP and Matthay KK. Neuroblastoma: clinical and biological approach to risk stratification and treatment. *Cell Tissue Res.* (2018) 1-15.
- [11] Group UCSW. United States cancer statistics: 1999–2006 incidence and mortality web-based report. Atlanta: US Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. <http://cdc.gov/uscs> (2010).
- [12] Ouyang L, Shi Z, Zhao S, Wang FT, Zhou TT, Liu B and Bao JK. Programmed cell death pathways in cancer: a review of apoptosis, autophagy and programmed necrosis. *CELL PROLIFERAT.* (2012) 45(6): 487-498.
- [13] Bai L and Wang S. Targeting apoptosis pathways for new cancer therapeutics. *Annu. Rev. Med.* (2014) 139-155.

- [14] Xu X, Zhang Y, Qu D, Jiang T and Li S. Osthole induces G2/M arrest and apoptosis in lung cancer A549 cells by modulating PI3K/Akt pathway. *J. Exp. Clin. Cancer. Res.* (2011) 30 (1): 33.
- [15] Zhou X, Wang X-B, Wang T and Kong L-Y. Design, synthesis, and acetylcholinesterase inhibitory activity of novel coumarin analogues. *Bioorg. Med. Chem.* (2008) 16(17): 8011-8021.
- [16] Zahir NV, Nakhjavani M, Hajian P, Shirazi FH and Mirzaei H. Evaluation of Silibinin effects on the Viability of HepG2 (Human hepatocellular liver carcinoma) and HUVEC (Human Umbilical Vein Endothelial) cell lines. *Iran J. Pharm. Res.* (2018) 17(1): 261
- [17] Vakili N, Nakhjavani M, Mirzayi H and Shirazi FH. Studying silibinin effect on human endothelial and hepatocarcinoma cell lines. *Res. Pharm. Sci.* (2012) 7(5): 174.
- [18] Lacy A and O'Kennedy R. Studies on coumarins and coumarin-related compounds to determine their therapeutic role in the treatment of cancer. *Curr. Pharm. Des.* (2004) 10(30): 3797-3811.
- [19] Jain P and Joshi H. Coumarin: Chemical and pharmacological profile. *J. Appl. Pharm. Sci.* (2012): 236-240.
- [20] Cooke D, Fitzpatrick B, O'Kennedy R, Mc Cormack T and Egan D. Coumarin Biochemical Profile and Recent Developments. *J.WS.* (1997).
- [21] Cooke D. Recent Advances on Coumarin and its Synthetic Derivatives. *D.C.U.* (1999) 12: 125-139.
- [22] Lee S-J, Lee U-S, Kim W-J and Moon S-K. Inhibitory effect of esculetin on migration, invasion and matrix metalloproteinase-9 expression in TNF- α -induced vascular smooth muscle cells. *MOL. MED. REP.* (2011) 4 (2): 337-341.
- [23] Nitiema LW, Savadogo A, Simporé J, Dianou D and Traore AS. In vitro antimicrobial activity of some phenolic compounds (coumarin and quercetin) against gastroenteritis bacterial strains. *Int. J. Microbiol. Res.* (2012) 3 (3): 183-187.
- [24] Ebbinghaus S, Mahler J and Marshall M. Novel Coumarin activity on albino rats and shows anti-coagulant and anti-oxidant activity. *J.W.S.* (1997) 45: 195-209.
- [25] Huang XY, Shan ZJ, Zhai HL, Su L and Zhang XY. Study on the anticancer activity of coumarin derivatives by molecular modeling. *Chem. Biol. Drug. Des.* (2011);78(4): 651-658.
- [26] Yeh JY, Coumar MS, Horng JT, Shiao HY, Kuo FM, Lee HL, Chen IC, Chang CW, Tang WF, Tseng SN and Chen CJ. Anti-Influenza Drug Discovery: Structure– Activity Relationship and Mechanistic Insight into Novel Angelicin Derivatives. *J. Med. Chem.* (2010);53(4): 1519-1533.
- [27] Asif M. Pharmacologically potentials of different substituted coumarin derivatives. *Chem. Int.* (2015);1(1): 1-11.
- [28] Iranshahi M, Arfa P, Ramezani M, Jaafari MR, Sadeghian H, Bassarello C, Piacente S and Pizza C. Sesquiterpene coumarins from *Ferula szowitsiana* and in vitro antileishmanial activity of 7-prenyloxycoumarins against promastigotes. *Phytochemistry* (2007);68(4): 554-561.
- [29] Russo P, Cimoli G, Valenti M, Sessa FD, Parodi S and Pommier Y. Induction of DNA double-strand breaks by 8-methoxycaffeine: cell cycle dependence and comparison with topoisomerase II inhibitors. *Carcinogenesis* (1994);15(11): 2491-2496.
- [30] Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. IMMUNOL. METHODS* (1983);65(1): 55-63.
- [31] Zhuang C, Itoh H, Mizuno T and Ito H. Antitumor active fucoidan from the brown seaweed, *umitoranoo* (*Sargassum thunbergii*). *Biosci. Biotechnol. Biochem.* (1995);59(4): 563-567.
- [32] Tayarani-Najaran Z, Mousavi SH, Vahdati-Mashhadian N, Emami SA and Parsaee H. *Scutellaria litwinowii* induces apoptosis through both extrinsic and intrinsic apoptotic pathways in human promyelocytic leukemia cells. *Nutr. Cancer* (2012) 64(1): 80-88
- [33] Singh RK, Lange TS, Kim KK and Brard L. A coumarin derivative (RKS262) inhibits cell-cycle

- progression, causes pro-apoptotic signaling and cytotoxicity in ovarian cancer cells. *Invest. New Drugs* (2011);29(1): 63-72.
- [34] Musa MA, Joseph MY, Latinwo LM, Badisa V and Cooperwood JS. In Vitro Evaluation of 3-Arylcoumarin Derivatives in A549 Cell Line. *Anticancer Res.* (2015);35(2): 653-659.
- [35] Khaghanzadeh N, Mojtahedi Z, Ramezani M, Erfani N and Ghaderi A. Umbelliprenin is cytotoxic against QU-DB large cell lung cancer cell line but anti-proliferative against A549 adenocarcinoma cells. *DARU* (2012);20(1): 69.
- [36] Wang J, Lu M, Dai H, Zhang S, Wang H and Wei N. Esculetin, a coumarin derivative, exerts in vitro and in vivo antiproliferative activity against hepatocellular carcinoma by initiating a mitochondrial-dependent apoptosis pathway. *Braz. J. Med. Biol. Res.* (2015);48(3): 245-253.
- [37] Zhang W, Li Z, Zhou M, Wu F, Hou X, Luo H, Liu H, Han X, Yan G, Ding Z and Li R. Synthesis and biological evaluation of 4-(1, 2, 3-triazol-1-yl) coumarin derivatives as potential antitumor agents. *Bioorg. Med. Chem. Lett.* (2014);24(3): 799-807.
- [38] Hoult J and Paya M. Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen. Pharmacol.* (1996);27(4): 713-722.
- [39] Klenkar J and Molnar M. Natural and synthetic coumarins as potential anticancer agents. *J. Chem. Pharm. Res.* (2015);7(7): 1223-1238.
- [40] Jun M, Bacay AF, Moyer J, Webb A and Carrico-Moniz D. Synthesis and biological evaluation of isoprenylated coumarins as potential anti-pancreatic cancer agents. *Bioorg. Med. Chem. Lett.* (2014);24(19): 4654-4658.
- [41] Bourgaud F, Hehn A, Larbat R, Doerper S, Gontier E, Kellner S and Matern U. Biosynthesis of coumarins in plants: a major pathway still to be unravelled for cytochrome P450 enzymes. *Phytochem Rev.*(2006);5(2-3): 293-308.
- [42] Farooq S. 555 medicinal plants. Field and laboratory manual (identification with its phytochemical and in vitro studies data): International Book Distributors; India (2005).
- [43] Rosselli S, Maggio AM, Faraone N, Spadaro V, Morris-Natschke SL, Bastow KF, Lee KH and Bruno M. The cytotoxic properties of natural coumarins isolated from roots of *Ferulago campestris* (Apiaceae) and of synthetic ester derivatives of aegelinol. *Nat. Prod. Commun.* (2009) 4(12): 1701-1706.
- [44] Crombie L, Games D and McCormick A. Isolation and structure of mammea A/BA, A/AB and A/BB: A group of 4-aryl-coumarin extractives of *mammea americana* L. *Tetrahedron Lett.* (1966)7(2): 145-149.
- [45] Poole SK and Poole CF. Thin-layer chromatographic method for the determination of the principal polar aromatic flavour compounds of the cinnamons of commerce. *A.N.A.L.* (1994);119(1): 113-120.
- [46] Soltani F, Mosaffa F, Iranshahi M, Karimi G, Malekaneh M, Haghghi F and Behravan J. Auraptene from *Ferula szowitsiana* protects human peripheral lymphocytes against oxidative stress. *Phytother Res.* (2010) 24(1): 85-89.
- [47] Lariche N, Mesbah L, Benguedouar L and Zelligui A. Ferulenol, a Sesquiterpene Coumarin, Induce Apoptosis via Mitochondrial Dysregulation in Lung Cancer Induced by Benzo [a] pyrene: Involvement of Bcl2 Protein. *Anticancer Agents. Med. Chem.* (2017): 1357-1362.
- [48] Salla S, Sunkara R, Walker LT and Verghese M. Antioxidant and Apoptotic Activity of Papaya Peel Extracts in HepG2 Cells. *Food Nutr. Sci.* (2016) 7 (6): 485.
- [49] Egan D, James P, Cooke D and O'Kennedy R. Studies on the cytostatic and cytotoxic effects and mode of action of 8-nitro-7-hydroxycoumarin. *Cancer Lett.* (1997) 118 (2): 201-211.
- [50] Brohem CA, Sawada TC, Massaro RR, Almeida RL, Rivelli DP, Ropke CD, Da Silva VV, De Lima TM, Curi R, Barros SB, Maria-Engler SS. Apoptosis induction by 4-nerolidylcatechol in melanoma cell lines. *Toxicol. In Vitro* (2009) 23(1): 111-119.

[51] Borhani N, Manoochehri M, Gargari SS, Novin MG, Mansouri A and Omrani MD. Decreased expression of proapoptotic genes caspase-8-and

BCL2-associated agonist of cell death (BAD) in ovarian cancer. *Clin. Ovarian Cancer Other Gynecol. cancer* (2014): 18-23.